

For the use of a Registered Medical Practitioner or a Hospital only.

**Human Albumin I.P. 20% Solution
(Xtra Purified)**

**ALBUCEL XP®
For Intravenous Use Only**

DESCRIPTION

ALBUCEL XP® is albumin (Human) 20%, prepared from the pools of adult human venous plasma obtained from healthy donors. AlbuCEL XP® is a sterile, liquid preparation of albumin in aqueous solution. 20 % of human albumin solution which is osmotically equivalent to 400 mL of normal human plasma.

PRODUCT SAFETY

Collected blood plasma which used in manufacturing of ALBUCEL XP®, screened for the mandatory infectious diseases. Only on being declared negative to HBsAg, HIV I & II antibodies, HCV RNA and antibodies against HCV the plasma is used for processing.

The manufacturing procedure incorporates Heat Pasteurization (60°C for 10 hours), which inactivates viruses. After manufacturing, the product is tested as per specification and indicates the product is non-reactive from viruses like HIV, HBsAg, HCV. Multiple steps have been applied to product safety assurance; there is a very remote probability that unknown infectious agents may be present in these products like newer emerging viruses and theoretical CJD (Creutzfeldt Jakob Disease). The process parameters, characterizations and final product quality are designed such, that they meet the regulatory requirements. Records of blood donors whose plasma have been used for manufacturing of this product have been maintained for at least ten years at the site of origin. AlbuCEL XP® contains no preservative and is free from plasma protein associated with the blood clotting mechanism and blood group antibodies. **Abbreviation:** HIV: Human Immunodeficiency Virus; HCV: Hepatitis C Virus; HBsAg: Hepatitis B surface antigen.

COMPOSITION

Each vial contains:

Concentration Available	20%
Pack Size	50ml and 100ml
Total Protein	200 g/L
Sodium Caprylate (as stabilizer)	6.648 g/L
Na ⁺ Contents	Not more than 160 mmol/L
K ⁺ Contents	Not more than 2 mmol/L
Aluminum Contents	Not more than 200 µg/L

Does not contain preservative.

CLINICAL PHARMACOLOGY

No pharmacokinetic or pharmacodynamic studies with AlbuCEL XP (Human Albumin) have been conducted.

Mechanism of Action

Albumin is responsible for 75-80% of the colloid osmotic pressure of normal plasma. Albumin stabilizes circulating blood volume and is a carrier of hormones, enzymes, medical products and toxins. Albumin is a protein with a total extravascular mass of approximately 160 g and an intravascular mass of about 120 g.

NON-CLINICAL TOXICOLOGY

No non-clinical toxicology studies with Albumin (Human) have been conducted. Human Albumin is a normal constituent of human plasma and acts like physiological albumin.

INDICATIONS AND USAGE

Hypovolemic Shock

AlbuCEL XP® is hyperoncotic solution. AlbuCEL XP® is indicated in the emergency treatment of hypovolemia with or without shock associated with blood loss, trauma and surgical procedure. On intravenous infusion, it will expand the plasma volume by an additional amount, three to four times the volume actually administered, by withdrawing fluid from the interstitial spaces, provided the patient is normally hydrated interstitially or there is interstitial edema. If the patient is dehydrated, additional crystalloids must be given. The patient's hemodynamic response should always be monitored with usual precautions against circulatory overload. If there has been considerable loss of red blood cells, transfusion with packed red blood cells is indicated. The total dose should not exceed the level of albumin found in the normal individual; i.e., about 2 g per kg body weight in the absence of active bleeding. AlbuCEL XP®, with appropriate crystalloids, may offer therapeutic advantages in oncotic deficits or in long-standing shock where treatment has been delayed.

Burn Therapy

An optimal therapeutic regimen with respect to the administration of colloids, crystalloids, and water following extensive burns has not been established. During the first 24 hours after sustaining thermal injury, large volumes of crystalloids are infused to restore the depleted extracellular fluid volume. Beyond 24 hours, AlbuCEL XP® is used in conjunction with adequate infusions of crystalloid to counteract hemoconcentration and the hypoproteinemia, electrolytes and water that usually follow severe burns.

Hypoproteinemia (With or Without Edema)

Albumin is indicated in the treatment of hypoproteinemia caused by a loss of plasma protein. Loss of plasma protein may occur through decreased absorption in gastrointestinal disorder, inadequate synthesis in chronic liver diseases or excessive urinary catabolism in chronic liver diseases. Albumin serves to restore colloidal osmotic pressure and in, conjugation with a diuretic, promotes diuresis. During major surgery, patients can lose over half of their circulating albumin with the attendant complications of oncotic deficit. A similar situation can occur in sepsis or intensive care patients. Treatment with AlbuCEL XP® may be of value in such cases.

Ascites

Removal of ascitic fluid from a patient with cirrhosis may cause changes in cardiovascular function and even result in hypovolemic shock. In such circumstances, albumin infusion is used to support the blood volume.

Renal Dialysis

Although not part of the regular regimen of renal dialysis, AlbuCEL XP® can be used as an adjunct in patients who are undergoing long term hemodialysis and are susceptible to shock or hypotension, or in dialysis patient who are hypovolemic and may not tolerate large volumes of crystalloids infusion as treatment for shock and hypovolemia. The usual volume administered is about 100 mL, taking particular care to avoid fluid overload as these patients are often fluid overloaded and cannot tolerate substantial volumes of salt solution. AlbuCEL XP® solution is suitable for use in dialysis patients, as aluminum content is less than 200 µg.

Acute Liver Failure

In the uncommon situation of rapid loss of liver function with or without coma, administration of AlbuCEL XP® may serve the double purpose of supporting the colloid osmotic pressure of the plasma as well as binding excess plasma bilirubin

Neonatal Hemolytic Disease

The administration of AlbuCEL XP® may be indicated prior to exchange transfusion, in order to bind free bilirubin, thus lessening the risk of kernicterus. AlbuCEL XP® solution is suitable for use in premature babies, as aluminum content is less than 200 µg.

DOSAGE AND ADMINISTRATION

AlbuCEL XP® should always be administered intravenously. The total dosage will vary with the individual. AlbuCEL XP® may be administered either undiluted or diluted in 0.9% Sodium Chloride solution (normal saline) or 5% Dextrose in Water. Whenever dilution of albumin human is necessary, the oncotic and osmotic properties as well as the tonicity of the resultant dilution must be considered. If sodium restriction is required, AlbuCEL XP® should only be administered either undiluted or diluted in a sodium-free carbohydrate solution such as 5 % Dextrose in Water. Because of risk of potentially life-threatening hemolysis and acute renal failure, albumin human must not be diluted with sterile water.

The concentration of the albumin preparation, dosage and the infusion-rate should be adjusted to the patient's individual requirements. The dose required depends on the size of the patient, the severity of trauma or illness and on continuing fluid and protein losses.

If AlbuCEL XP® is to be administered, hemodynamic performance should be monitored regularly; this may include:

- arterial blood pressure and pulse rate
- central venous pressure
- pulmonary artery wedge pressure
- urine output
- electrolyte
- hematocrit/hemoglobin

Measures of adequacy of circulating volume and not plasma albumin levels should be used to determine the dose required. If large volumes are administered, the product should be warmed to room temperature before use. In the treatment of shock, an initial dose of 100 mL of the 20% albumin solution is given as rapidly as tolerated. If response within 30 minutes is inadequate, an additional 100 mL of 20% albumin solution may be given. In the patient with a slightly low or normal blood volume, the rate of administration should be 1 mL per minute. If more than 250 mL are given, or if hemorrhage has occurred, the administration of packed red blood cells may be desirable.

In severe burns, immediate therapy should include large volumes of crystalloid with lesser amounts of 20% albumin solution to maintain an adequate plasma volume and protein content. After the first 24 hours, the ratio of albumin to crystalloid may be increased to establish and maintain a plasma albumin level of about 2.5 g/100 mL or a total serum protein level of about 5.2 g/100 mL. In acute hypoproteinemia 250-350 mL of 20% albumin may be required to reduce edema and to bring serum protein values to normal. Since such patients usually have approximately normal blood volume, the rate of administration should not be greater than 3 ml per minute to avoid circulatory embarrassment.

The initial dosage in children will vary with the clinical state and body weight. A dose one-quarter to one-half the adult dose may be administered, or dosage may be calculated on the basis of 1 - 3 mL of Albumin 20% per kg of body weight. The usual rate of administration in children should be one-quarter the adult rate. For infants suffering from hemolytic disease of the newborn the appropriate dose for binding of free serum bilirubin is 1 gram per kilogram of body weight. This may be administered before or during the exchange procedure. Caution must be observed in hypervolemic infants.

Parenteral drug products should be inspected visually for particulate matter and discoloration prior to administration.

USE IN SPECIAL POPULATIONS

Pregnancy

Pregnancy Category C. Animal reproduction studies have not been conducted with AlbuCEL XP®. It is also not known whether AlbuCEL® can cause fetal harm when administered to a pregnant woman or can affect reproductive capacity. AlbuCEL XP® should be given to a pregnant woman only if clearly needed.

Pediatric Use

No clinical studies establishing safety and effectiveness of albumin have been conducted in pediatric patients. However, extensive experience in patients suggests that children respond to albumin in the same manner as adults. There are no known adverse reports of human albumin usage in children, if dose is appropriate for body weight of child.

CONTRAINDICATIONS

Human albumin is contraindicated in certain patients e.g. those with a history of congestive cardiac failure, renal insufficiency or stabilize chronic anemia are at special risk of developing circulatory overload. Human albumin is contraindicated in patient with hypersensitivity to albumin preparations or to any of the excipients present in the solution.

WARNINGS

AlbuCEL XP® is made from human plasma. Products made from human plasma may contain infectious agents, such as viruses, and, theoretically, the Creutzfeldt - Jakob Disease (CJD) agent that can cause disease. The theoretical risk for transmission of CJD is considered extremely remote. No cases of transmission of viral diseases or CJD have ever been identified for albumin. The risk that such products will transmit an infectious agent has been reduced by screening plasma donors for prior exposure to certain viruses, by testing for the presence of certain current virus infections, and by inactivating and/or removing certain viruses. The manufacturing procedure for AlbuCEL XP® includes processing steps designed to reduce further the risk of viral transmission. AlbuCEL XP® is pasteurized in the final container at 60.0 °C for 10 hours. Virus eliminated /inactivation is also achieved by cold alcohol fractionation process. Despite these measures, such products can still potentially transmit disease. There is also the possibility that unknown infectious agents may be present in such products. Individuals who receive infusions of blood or plasma products may develop signs and/or symptoms of some viral infections, particularly hepatitis C. All infections thought by a physician possibly to have been transmitted by this product should be reported by the physician or other healthcare provider. The physician should discuss the risks and benefits of this product with the patient, before prescribing or administering it to the patient. As with any hyperoncotic protein solution likely to be administered in large volumes, severe hemolysis and acute renal failure may result from the inappropriate use of Sterile Water for Injection as a diluent for Albumin 20%. Acceptable diluents include 0.9% Sodium Chloride or 5% Dextrose in Water. Please refer to the **DOSAGE AND ADMINISTRATION** section for recommended diluents.

Bottles which are cracked or which have been previously entered or damaged should not be used, as this may have allowed the entry of microorganisms. AlbuCEL XP® contains no preservative.

OVERDOSE:

Hypervolaemia may occur if the dosage and rate of infusion are too high. At the first clinical signs of cardiovascular overload (headache, dyspnoea, jugular vein congestion), or increased blood pressure, raised central venous pressure and pulmonary oedema, the infusion should be stopped immediately and the patient's haemodynamic parameters carefully monitored.

PRECAUTIONS

Certain solutions containing protein hydrolysates or alcohol must not be infused through the same administration set in conjunction with albumin since these combinations may cause the proteins to precipitate.

Albumin should be administered with caution to patients with low cardiac reserve.

Albumin should be used with caution in patients who are at increased risk of developing circulatory overload and its consequences or hemodilution could represent a special risk for the patient. Examples of such conditions are:

- Hypertension
- Esophageal varices
- Pulmonary edema Hemorrhagic diathesis
- Renal and post-renal anuria

Rapid infusion may cause vascular overload with resultant pulmonary edema. Patients should be closely monitored for signs of increased venous pressure. At the first clinical signs of cardiovascular overload (headache, dyspnea, jugular vein congestion), or increased blood pressure, raised venous pressure and pulmonary edema, the infusion is to be stopped immediately.

A rapid rise in blood pressure following infusion necessitates careful observation of injured or postoperative patients to detect and treat severed blood vessels that may have bled at a lower pressure.

Patients with marked dehydration require administration of additional fluids. Albumin may be administered with the usual dextrose and saline intravenous solutions.

20 % *albumin* solution is relatively low in electrolytes compared to the 5% albumin solution. When albumin is given the electrolyte status of the patient should be monitored and appropriate steps taken to restore or maintain the electrolyte balance.

If comparatively large volumes are to be replaced, controls of coagulation and hematocrit are necessary. Care must be taken to ensure adequate substitution of other blood constituents (coagulation factors, electrolytes, platelets and erythrocytes).

DRUG INTERACTIONS

AlbuCEL XP® must not be mixed with other medicinal product, whole blood, packed red cells, as well as the standard carbohydrate and electrolyte solutions intended for intravenous use. It should, however, not be mixed with protein hydrolysates, amino acid solutions nor those containing alcohol. Components used in the packaging of AlbuCEL XP® 20% are latex-free.

ADVERSE REACTIONS

Adverse reactions with albumin are rare. These reactions normally disappear rapidly when the infusion rate is slowed down or the infusion is stopped. In case of severe reactions, the infusion should be stopped and an appropriate treatment should be initiated. If administration has been stopped and the patient requires additional AlbuCEL XP®, material from a different lot should be used.

Adverse reactions include:

- Allergic or pyrogenic reaction: anaphylaxis, which may be severe, and hypersensitivity reactions including pyrexia, chills, urticaria, skin rash, pruritus, edema, erythema, hypotension and bronchospasm.
- Psychiatric: confusional state
- Nervous system: headache
- Cardiac: tachycardia, cardiac failure
- Vascular: hypotension, hypertension, flushing
- Respiratory: dyspnoea
- Gastrointestinal: nausea, vomiting
- Skin and subcutaneous tissue: erythematous, hyperhidrosis

HOW SUPPLIED

AlbuCEL XP® is available as 20 % intravenous infusion single dose hermetic container, containing 20 g human albumin per 50 ml and 100ml.

STORAGE

Store below 25° C

Do not begin administer if the container has been opened for more than 4 hours. Any remnant portions to be discarded.

Do not use if the solution is turbid or any particulate matters observed.

Keep the container in the outer carton in order to protect from light.

EXPIRY

36 months from the date of manufacture. Do not use after expiry date.

Report suspected adverse reaction at: Hemofluidsafety@intaspharma.com

Date of preparation : 29/09/23

Manufactured and Marketed by: 

INTAS PHARMACEUTICALS LTD.

Plot No. 496/1/A&B, Sarkhej-Bavia Highway,
Village: Matoda, Taluka: Sanand,
Ahmedabad-382213, Gujarat, (INDIA)