

130 mm

290 mm

For the use of a Registered Medical Practitioner only.
Human Normal Immunoglobulin for Intravenous Use I.P. 10% Solution
GLOBUCEL® VF-10
ग्लोबुसेल® VF-10

Manufactured from human plasma - Virus Filtered - For Intravenous Use Only

GENERIC NAME

Human normal immunoglobulin for intravenous use I.P. 10% solution (Virus Filtered)

QUALITATIVE AND QUANTITATIVE COMPOSITION

Concentration Available: 10%
Pack size: 50 mL and 100 mL
Each vial contains:
Total Protein: 100 g/L
Immunoglobulin G: ≥ 95%
Maltose (as stabilizer): 100 g/L
pH Content: 7.0-7.5
IgG subclass: Normal Distribution
Water for Injection: q.s.

DOSEAGE FORM AND STRENGTH

Human normal immunoglobulin solution for intravenous (I.V.) use only.
It is supplied as 50 mL (5 g) and 100 mL (10 g) vial.

CLINICAL PARTICULARS

THERAPEUTIC INDICATION

Immunoglobulin preparations are indicated in several clinical conditions. An approved list of clinical conditions where IVIG is indicated is as under:

Primary Immunodeficiency (PID) Syndromes

Replacement therapy to promote passive immunity: the following PID syndromes can be treated with intravenous replacement of IgG and are considered well established:

- Congenital agammaglobulinemia and hypogammaglobulinemia
- Common variable immunodeficiency
- X-linked agammaglobulinemia
- Wiskott-Aldrich syndrome
- Severe combined immunodeficiencies.

IVIG may be preferred in patients who require an immediate or large increase in intravascular immunoglobulin concentrations, in patients with small muscle mass, and in patients with bleeding tendencies in whom Intramuscular (IM) injections are contraindicated.

Secondary Immunodeficiency (SID) Syndromes

IgG can also be used as replacement therapy in:

- Secondary hypogammaglobulinemia in patients with B-cell chronic lymphocytic leukaemia (CLL) or multiple myeloma (MM) with recurrent infections.
- Paediatric HIV-1 infection who have bacterial infections: HIV-infected infants and children with hypogammaglobulinemia (IgG <400 mg/dL) should receive IVIG (400 mg/kg once every 2-4 weeks) to prevent serious bacterial infections.

Kawasaki Syndrome

IVIG is used in conjunction with aspirin therapy for initial treatment of the acute phase of Kawasaki disease. Approximately 21% of patients with Kawasaki disease fail to respond to initial treatment with IVIG and aspirin therapy and have persistent fever or recurrent fever after an initial afebrile period. Retreatment with IVIG (within 10-14 hours of persistent or recrudescence fever) and continued aspirin therapy usually is recommended for these patients.

Idiopathic Thrombocytopenic Purpura

IVIG is indicated for the treatment of acute or chronic (e.g., >6 months duration) idiopathic thrombocytopenic Purpura.

Allogeneic Bone marrow transplantation (BMT)

In adults and children undergoing BMT, IVIG can be used to decrease the risk of infections (e.g., septicemia), interstitial pneumonia of infectious or idiopathic etiologies, and acute graft-versus-host disease.

G Guillain-Barre syndrome

IVIG initiated within 2 weeks of symptom onset appears to be as effective as plasma exchange.

Chronic Inflammatory Demyelinating Polyneuropathy (CIDP)

IVIG may be used for the treatment of chronic inflammatory demyelinating polyneuropathy to improve neuromuscular disability and impairment and for maintenance therapy to prevent relapse.

Multifocal Motor Neuropathy (MMN)

PHARMACOLOGY AND METHOD OF ADMINISTRATION

For intravenous use only

IVIG liquid should be inspected visually for particulate matter and discoloration prior to administration, whenever solution and container permit. Do not use if turbid and/or discoloration is observed. IVIG liquid must not be mixed with other medicinal products or administered simultaneously with other intravenous preparation in the same infusion set. Do not mix with intravenous immunoglobulin product containing other antibodies.

IVIG liquid should be at room temperature during administration. Any bottle that has been opened should be promptly. Partially used bottles should be discarded since this drug does not contain any preservatives.

Prior to initiation of IVIG infusion, ensure that patients are not volume depleted and are adequately hydrated. Individualize the rate of infusion based on the preparation and individual patient requirements.

In general, in patients receiving initial doses of IVIG or switching from one IVIG preparation to another, initiate the infusion rate at the lower end of the recommended range and increase to the maximum recommended rate only after the patient has tolerated several infusions at an intermediate infusion rate.

If an adverse reaction occurs during the IVIG infusion, decrease the rate of infusion or stop the infusion until the reaction subsides. Do not administer by rapid IV infusion in patients with or at risk for renal dysfunction or thrombotic events.

Risk factors should be identified, such as pre-existing renal insufficiency, diabetes mellitus, hypovolemia, overweight, concomitant nephrotoxic medications, or over the age of 65. Assume that patients are not volume depleted prior to the initiation of the infusion of IVIG. Patients should be observed for at least 20 minutes after administration.

Respective of blood group, it can be transferred to all recipients. Primary Immunodeficiency Syndrome

As there are significant differences in the half-life of IgG among patients with primary humoral immunodeficiencies, the frequency and amount of immunoglobulin therapy may vary from patient to patient. The proper amount can be determined by monitoring clinical response. The dosage regimen should achieve a trough level of IgG (measured before the next infusion) of at least 4-6 g/L. The dose of IVIG liquid for replacement therapy in primary humoral immunodeficiency diseases is 300 to 600 mg/kg body weight (8-12 mL/kg) administered every 3 to 4 weeks. The dosage may be adjusted over time to achieve the desired trough levels and clinical responses. Three to six months are required after the initiation of therapy for equilibration to occur.

Kawasaki Disease

For initial treatment of acute phase, AHA, AAP, and ACCP recommend a single dose of 2 g/kg of IVIG given in conjunction with aspirin (80-100 mg/kg daily for up to 14 days, then 1-5 mg/kg once daily for 6-8 weeks); initiate as soon as possible (optimally within 7-10 days of disease onset). If there is no response (i.e., fever persists or recurs ≥36 hours after initial IVIG dose), retreatment with another single dose of 2 g/kg of IVIG (given within 24-48 hours of persistent or recrudescence fever) and continued aspirin therapy is recommended.

Idiopathic Thrombocytopenic Purpura (ITP)

For initial treatment of acute phase, 200-400 mg/kg once daily for 5 consecutive days. In acute childhood ITP, if an initial platelet count response to the first 2 doses is adequate (30,000-50,000/mm3), discontinue therapy after the second day of the 5-day regimen. For treatment of chronic ITP, if platelet count decreases to <100,000/mm3 and/or clinically important bleeding becomes apparent following initial induction therapy, administer 400 mg/kg as a single maintenance infusion. If an adequate response does not occur, increase the maintenance dose to 800-1000 mg/kg given as a single infusion.

Prevention of Serious Infections in HIV-Infected Individuals

Infants and children with hypogammaglobulinemia (IgG <400 mg/dL): ACIP, AAP, CDC, NIH, and other experts recommend 400 mg/kg of IVIG once every 2-4 weeks.

G Guillain-Barre Syndrome

European Federation of Neurological Societies (EFNS) and others recommend 0.4 g/kg daily for 5 days. The dosage regimens given below are general guidelines. Actual dose regimen will be decided base on physician's decision of patient's clinical condition.

Multifocal Motor Neuropathy (MMN)

Starting dose: 2 g/kg given over 2-5 days. Maintenance dose: 1 g/kg every 2 to 4 weeks or 2 g/kg every 4 to 8 weeks.

First Infusion:

It is recommended that GlobuceL® VF-10 liquid be initially infused at infusion rates stated below, at least until the physician has had adequate experience with a given patient.

Table with 4 columns: Steps, Time, First infusion of GlobuceL® VF-10 Rate (ml/kg/hr), and Subsequent Infusions: Steps, Time, Subsequent infusion of GlobuceL® VF-10 Rate (ml/kg/hr). It details the infusion schedule for the first and subsequent infusions.

*Rate not to exceed 300 ml/hr. For patients judged to be at risk for developing renal dysfunction or thrombotic events, administer IVIG liquid at the minimum infusion rate practicable. Data are not available to date to identify maximum safe dose, concentration, and rate of infusion in patients at risk for renal dysfunction. Ensure that patients with pre-existing renal insufficiency are not volume depleted; discontinue IVIG if renal function deteriorates.

Doorage recommendation for human intravenous immunoglobulin

Table with 2 columns: No. Indication and Dose. It lists various clinical conditions and the corresponding IVIG dosage recommendations.

USE IN SPECIAL POPULATION

Pregnancy

US-FDA Pregnancy category C. Animal reproduction studies have not been conducted with intravenous immunoglobulin (IVIG). It is also not known whether intravenous immunoglobulin can cause fetal harm when administered to a pregnant woman or can affect reproduction capacity. There is the possibility of parvovirus B19 infection due to administration of this drug. In case of infection, fetal disorder such as abortion, fetal hydrops, and fetal death may occur. Intravenous immunoglobulin should be given to a pregnant woman only if clearly needed.

Nursing Mothers

Intravenous immunoglobulin has not been evaluated in nursing mothers.

Pediatric Use

Safety and efficacy of intravenous immunoglobulin have not been established in children under 2 years of age. Intravenous immunoglobulin was evaluated in 11 pediatric subjects (age range 6- 16 years). There were no obvious differences observed between adults and pediatric subjects with respect to pharmacokinetics, efficacy and safety. No pediatric specific dose requirements were necessary to achieve the desired serum IgG levels.

Geriatric Use

Patients > 65 years of age may be at increased risk for developing certain adverse reactions such as thrombotic events and acute renal failure (See Warnings and Precautions). In the clinical trial only 4 geriatric patients (> 65 years) were enrolled, a number insufficient to determine whether geriatric patients respond differently from younger subjects. In these 4 patients no particular issue was observed.

CONTRAINDICATIONS

IVIG is contraindicated in individuals who have had anaphylactic or severe systemic reactions to immunoglobulin or any ingredients in the formulation. Epinephrine should be available for immediate treatment of an anaphylactic reaction if it occurs.

IVIG is contraindicated in individuals with selective IgA deficiency or IgA deficiency with antibodies against IgA, since these individuals may have antibodies to IgA (or develop antibodies following administration of IVIG) or other blood products containing IgA.

Acute hypersensitivity reaction to corn; this product contains maltose derived from corn.

SPECIAL WARNINGS AND PRECAUTIONS FOR USE

Human normal immunoglobulin solution is human plasma-derived immunoglobulin preparation. Bring the medicine to room or body temperature before use.

Sensitivity

Severe hypersensitivity reactions, including anaphylaxis, reported rarely following administration of Intravenous Immunoglobulin (IVIG), Intramuscular Immunoglobulin (IMIG) or Subcutaneous Immunoglobulin (SCIG). Epinephrine and antihistamines should be readily available in case anaphylaxis or an anaphylactoid reaction occurs. If a severe hypersensitivity reaction occurs, discontinue immune globulin immediately and institute appropriate therapy as indicated. IgA deficient patients with antibodies against IgA are at greater risk of developing severe hypersensitivity and anaphylactoid reactions when administered IVIG (See CONTRAINDICATIONS). Patients known to have corn allergies should avoid using IVIG (See CONTRAINDICATIONS).

Infection Reactions

There is a risk of reactions including fever, chills, nausea, and vomiting upon IV infusion. These reactions generally appear 30 minutes to 1 hour after initiation of the infusion and include flushing of the face, tightness in the chest, chills, fever, dizziness, nausea, vomiting, diaphoresis, and hypotension or hypertension. Closely monitor for adverse reactions throughout the infusion since these reactions may rarely lead to shock.

Human normal immunoglobulin solution may cause a precipitous fall in BP and clinical manifestations of anaphylaxis, which appear to be related to the rate of infusion, do not exceed the recommended rate of infusion. If flushing, changes in BP or pulse, or other infusion reactions occur, slow or temporarily stop the infusion. In some cases when symptoms subside promptly, the infusion may be resumed at a rate that is comfortable for the patient. Stop infusion immediately if anaphylaxis or other severe reactions occur.

Acute renal failure

Renal dysfunction, acute renal failure, osmotic nephrosis, and death reported in patients receiving immune globulin. Patients at increased risk for acute renal failure include those with any degree of preexisting renal insufficiency, diabetes mellitus, volume depletion, sepsis, or paraproteinemia; those receiving concomitant nephrotoxic drugs; and/or those >65 years of age. To minimize risk of acute renal failure, ensure that patients are not volume depleted and are adequately hydrated prior to administration of IVIG. Always use lowest effective dosage at the minimum concentration available and at

the minimum practicable rate of infusion, especially in patients at increased risk for acute renal failure. Assess urine output and renal function including blood urea nitrogen (BUN) serum creatinine, prior to and at appropriate intervals during therapy with IVIG, especially in patients considered at increased risk for acute renal failure. If renal dysfunction occurs, consider discontinuing immune globulin therapy.

Risk of Transmissible Agents in Plasma-derived Preparations

Because immune globulin preparations are prepared from pooled human plasma, they may carry a risk of transmitting infectious agents, including the causative agents of viral hepatitis and HIV infection, and theoretically may carry a risk of transmitting the causative agent of Creutzfeldt-Jakob disease (CJD) or variant CJD (vCJD). Risk for transmission of recognized blood-borne viruses is considered to be low because plasma donors are screened for certain viruses (HBV, HCV, HIV) and viral reduction/inactivation procedures used in immune globulin production reduce the risk of transmission. Despite such stringent procedures, a risk of transmission still remains.

Because no purification method has been shown to be totally effective in removing the risk of viral infectivity from plasma-derived preparations and because new blood-borne viruses or other disease agents may emerge that may not be inactivated by the manufacturing process or various treatment procedures used, carefully weigh risk of pathogen transmission against the benefits of immune globulin therapy. Report all infections thought possibly to have been transmitted by immune globulin preparations to the appropriate manufacturer.

Thrombotic Effects

Thrombotic events (e.g., chest pain, MI, CHF, cerebral infarction, ischemic encephalopathy, severe headache requiring hospitalization, pulmonary embolism, retinal vein occlusion, peripheral venous thrombosis), including some fatalities, reported in patients receiving IVIG. IVIG-induced alterations of blood rheology (e.g., platelet activation, increased blood viscosity) and infusion-related hypertensive effects appear to contribute to the development of thrombotic complications. Patients with a history of atherosclerosis, multiple cardiovascular risk factors, hypertension, impaired cardiac output, cerebrovascular disease, coronary artery disease, coagulation or hypercoagulable disorders (e.g., factor V Leiden), prolonged periods of immobilization, advanced age, obesity, diabetes mellitus, acquired or inherited thrombotic disorder, previous thrombotic or thromboembolic event, or known or suspected hyperviscosity, and/or those receiving estrogen-containing products may be at increased risk. Weigh potential risks and benefits of immune globulin against those of alternative therapies in all patients in whom immune globulin is being considered.

Prior to immune globulin therapy, carefully evaluate patients with thrombotic risk factors (e.g., those with advanced age, hypertension, cerebrovascular disease, CAD, diabetes mellitus, high serum levels of a monoclonal protein, a history of prolonged immobilization [e.g., bed-bound], and/or a history of thrombotic episodes). Because of potential increased risk of thrombosis, consider baseline assessment of blood viscosity in patients at risk for hyperviscosity (e.g., those with cryoglobulins, fasting chylomicronemia/markedly high lipoproteins, or monoclonal gammopathies).

Hemolysis

IVIG and immune globulin subcutaneous can contain blood group antibodies which may act as hemolysins and induce in vivo coating of RBCs with immunoglobulin, causing a positive direct antiglobulin reaction and, rarely, hemolysis. Monitor for clinical signs and symptoms of hemolysis (e.g., increased heart rate, swelling, fatigue, difficulty breathing, yellowing of skin or eyes, dark-colored urine) and, if necessary, perform confirmatory laboratory testing. If a blood transfusion is indicated for a patient who developed hemolysis with clinically compromising anaemia after receiving immune globulin, ensure adequate cross-matching should be performed to avoid exacerbating on-going hemolysis.

Hemolytic anaemia also can develop subsequent to immune globulin therapy due to enhanced RBC sequestration and/or intravascular RBC destruction.

Transfusion-related Acute Lung Injury

Transfusion-related acute lung injury (noncardiogenic pulmonary edema) has been reported in patients receiving IVIG. Typically occurs within 1-6 hours after the infusion and is characterized by severe respiratory distress, pulmonary edema, hypoxemia, normal left ventricular function, and fever. Monitor patients receiving Human normal immunoglobulin solution for adverse pulmonary reactions. If transfusion-related acute lung injury is suspected, perform appropriate tests for the presence of antineutrophil antibodies and anti-human leukocyte antigen (HLA) antibodies in both the product and patient serum. Manage using oxygen therapy with adequate ventilatory support.

Aseptic Meningitis Syndrome

Aseptic meningitis syndrome reported infrequently in patients receiving immune globulin, especially at high doses (e.g., >1 g/kg) and/or by rapid IV infusion which should be rarely possible with Human normal immunoglobulin solution. Symptoms (e.g., severe headache, nuchal rigidity, drowsiness, fever, photophobia, painful eye movements, nausea, vomiting) may occur within several hours to 2 days following the administration.

Hyperproteinemia, Increased Viscosity, and Hyponatremia

Hyperproteinemia, increased serum viscosity, and hyponatremia may occur in patients receiving IVIG. The hyponatremia is likely to be pseudohyponatremia, as demonstrated by decreased calculated serum osmolality or elevated osmolal gap.

If hyponatremia occurs, it is critical to distinguish true hyponatremia from pseudohyponatremia. Treatment aimed at decreasing serum free water in patients with pseudohyponatremia may lead to volume depletion, a further increase in serum viscosity, and may predispose to thromboembolic events.

Volume Overload

High-dose IVIG regimens (1 g/kg daily for 1-2 days) used for treatment of chronic ITP are not recommended in individuals with expanded fluid volumes or when fluid volume may be a concern.

Blood Glucose Testing

IVIG preparations that contain maltose may cause falsely elevated results in blood glucose determinations with tests that use non-specific methods based on glucose dehydrogenase pyridoxal phosphate (GDH-PQQ) or glucose-dye oxidoreductase. This has resulted in the inappropriate administration of insulin, resulting in life-threatening hypoglycemia. Also, cases of true hypoglycemia may go untreated if the hypoglycemic state is masked by falsely elevated glucose readings. Accordingly, when administering IVIG, the measurement of blood glucose must be done with a glucose-specific method. The product information of the blood glucose testing system, including that of the test strips, should be carefully reviewed to determine if the system is appropriate for use with maltose-containing parenteral products.

DRUGS INTERACTIONS

Admixtures of intravenous immunoglobulin with other drugs and intravenous solutions have not been evaluated. It is recommended that intravenous immunoglobulin liquid be administered separately from other drugs or medications which the patient may be receiving. The product should not be mixed with IVIG from other manufacturers.

The infusion line may be flushed before and after administration of intravenous immunoglobulin with either normal saline or 5% dextrose in water. Various passively transferred antibodies in immunoglobulin preparations can confound the results of serological testing.

Antibodies in intravenous immunoglobulin (IVIG) may interfere with the response to live viral vaccines, including measles, mumps, and rubella virus vaccine live (MMR) and varicella virus vaccine live. No response to the interferon response to influenza virus vaccine live intranasal, yellow fever vaccine live, typhoid vaccine live oral, rotavirus vaccine live oral, zoster vaccine live oral, or poliovirus vaccine live oral. However, caution should be exercised during use of above vaccine and physicians should follow the prescribing information of respective vaccine. Physicians should be informed of recent therapy with IVIGs, so that administration of live viral vaccines, if indicated, can be appropriately delayed 3 or more months from the time of IVIG administration. In the case of measles, this impairment may persist for up to 1 year. These patients receiving live vaccines should have their antibody status checked.

EFFECTS ON ABILITY TO DRIVE AND USE MACHINES

No information on the ability to drive and use machines have been known. Patients may experience effects, such as dizziness or nausea, during treatment with Human normal immunoglobulin solution that might affect the ability to drive and use machines. If this happens, you should not drive or use machines until these effects have disappeared.

UNDESIRABLE EFFECTS

The most common adverse reactions reported in ≥ 5% of clinical trial subjects occurring during or within 48 hours of an infusion were headache, nausea, chills, asthenia (fatigue), pyrexia, upper respiratory tract infection, hyperhidrosis, and flushing. In post-marketing surveillance, serious adverse reactions reported with intravenous immunoglobulin were anaphylaxis, acute renal failure, myocardial infarction, cerebral vascular accident, transient ischemic attack, deep vein thrombosis, pulmonary embolism, aseptic meningitis, acute hemolysis, and TRALI. Rare cases of transient cutaneous reactions (including cutaneous lupus erythematosus) have been observed.

The following adverse reactions have been identified during post-approval use of IVIG products. Because these reactions are reported voluntarily from a population of uncertain size, it is not always possible to reliably estimate their frequency or to establish a causal relationship to IVIG products. Certain severe drug reactions may be related to the rate of infusion. Possible adverse reactions with human normal immunoglobulin solution are listed below.

Table listing adverse reactions categorized by system: Blood and lymphatic system disorders, Immune system disorders, Metabolic and nutritional disorders, Psychiatric disorders, Nervous system disorders, Cardiac system disorders, Vascular disorders, Respiratory, thoracic and mediastinal disorders, Gastrointestinal disorders, Skin and subcutaneous tissue disorders, Musculoskeletal and connective tissue disorders, Renal and urinary disorders, General disorders and administration site conditions, and Investigations.

OVERDOSE

Overdose is very unlikely to occur because Human normal immunoglobulin solution is usually administered under medical supervision. Overdose of Human normal immunoglobulin solution can lead to fluid overload and increased thickness of the blood (hyperviscosity), particularly in at-risk patients, including elderly patients or patients with impaired renal function. In case of overdose, further infusion should be halted and medical supervision is suggested, along with symptomatic management.

PHARMACOLOGICAL PROPERTIES

MECHANISM OF ACTION

Human normal immunoglobulin solution having very high titres of antibodies against the infectious agents. It will provide passive immunity for the prevention and treatment of the infections. The evidence and mechanism of action of IgG in different diseases has not been fully elucidated.

PHARMACODYNAMIC PROPERTIES

IVIG, human intravenous immunoglobulin, provides a broad spectrum of opsonic and neutralizing IgG antibodies against a wide variety of bacterial and viral agents reflecting the IgG activity found in the donor population. It has an IgG subclass distribution similar to that of native human plasma. IgG antibodies contained in IVIG provide passive immunity by increasing an individual's antibody titer and antigen-antibody reaction potential and prevent or modify certain infectious diseases in susceptible individuals. Adequate doses of IVIG can restore abnormally low IgG level to the normal range. The role of evidence and mechanism of action of IgG in different diseases has not been fully elucidated.

PHARMACOKINETIC PROPERTIES

Peak levels of IgG are reached immediately after infusion of IVIG in patients with primary immunodeficiency syndrome. Following infusion, IVIG products show a biphasic decay curve. The initial (α) phase is characterized by an immediate post-infusion peak in serum IgG and is followed by rapid decay due to equilibration between the plasma and extravascular fluid compartments until approximately half is partitioned in the extravascular space, equilibrium is reached between the intra and extravascular compartments. The second (β) phase is characterized by a slower and constant rate of decay. As a class, IgG survives longer in vivo than other serum proteins. Peak levels of IgG are reached within 30 minutes after an intravenous infusion of IVIG.

Half-life of IgG in individuals with normal serum IgG concentrations is around 18-25 days while it is 12-45 days in patients with immunodeficiencies. The half-life of IgG can vary considerably from person to person, however. In particular, high serum concentrations of IgG and hypermetabolism associated with fever and infection have been seen to coincide with a shortened half-life of IgG. IgG and IgG-complexes are broken down in cells of the reticuloendothelial system.

NONCLINICAL PROPERTIES

ANIMAL TOXICOLOGY OR PHARMACOLOGY

Being human plasma-derived proteins, safety testing in animals is not particularly relevant to correlate the safety of use in man. Moreover, as these human plasma proteins are more immunogenic to animals than to humans, toxicity and productivity of pre-clinical testing further diminish. In animals, single-dose toxicity testing is of little relevance and does not permit the evaluation of toxic and lethal doses or a dose-effect relationship. Repeated dose toxicity testing is impractical due to the development of antibodies to heterogeneous protein in animal models.

DESCRIPTION

Human Normal Immunoglobulin for Intravenous use 10% solution (Virus Filtered) is a sterile and solvent-detergent (S/D) treated preparation of highly purified Immunoglobulin G (IgG) intended for intravenous use. It is prepared from the large pools of the human or plasma obtained from the healthy donors. IVIG is used to provide passive immunity by increasing an individual's antibody titer and antigen-antibody reaction potential. IVIG also helps to prevent or modify certain infectious diseases in susceptible individuals.

PRODUCT SAFETY

Collected blood plasma which used in manufacturing of IVIG, screened for the mandatory infectious diseases. Only on being declared negative to HBsAg, HIV I & II antibodies, HCV RNA and antibodies against HCV the plasma is used for processing.

The manufacturing procedure incorporates two dedicated orthogonal virus clearance steps ensuring viral safety of the product. This includes solvent detergent treatment and virus retentive filtration. The use of 20 m virus retentive filter provides additional safety against enveloped and non-enveloped virus. The manufacturing process of GlobuceL® VF-10 shows significant viral reduction and inactivation, demonstrated by virus validation study. Below Table shows the virus clearance capacity of two steps during the manufacturing of GlobuceL® VF-10, expressed as the mean log10 reduction factor (RF).

Table showing virus clearance capacity of two steps during the manufacturing of GlobuceL® VF-10. Columns include Process, BVDV, HIV-1, PRV, HAV, and PPV. Rows include S/D Treatment, Virus Filtration (20N), and Cumulative minimum / mean reduction factor.

After manufacturing, the product is tested as per specification and that also indicates the product is non-reactive to viruses like HIV, HBsAg, HCV. Multiple tests have been applied to product safety assurance; there is a very remote probability that unknown infectious agents may be present in these products like newer emerging viruses and theoretical CJD (Creutzfeldt Jakob Disease). The process parameters, characterizations and final product quality are designed such, that they meet the regulatory requirements. Records of blood donors whose plasma have been used for manufacturing of this product have been maintained as per the applicable regulatory requirement at the site of origin.

Abbreviation: IVIG: Intravenous immunoglobulins; HIV: Human Immunodeficiency Virus; HCV: Hepatitis C Virus; HBsAg: Hepatitis B surface antigen.

PHARMACEUTICAL PARTICULARS

INCOMPATIBILITIES

In the absence of compatibility studies, this medicinal product must not be mixed with other medicinal products.

SHELF-LIFE

Two years from the date of manufacture. Do not use it after expiry.

PACKAGING INFORMATION

Container Closure: USP Type I glass vial with Brombutyl rubber stopper

IVIG is supplied as 10% solution containing 5g of human Normal Immunoglobulin per 50 ml and 10g of human Normal Immunoglobulin per 100 ml for intravenous administration.

STORAGE AND HANDLING INSTRUCTIONS

Store between +2°C to +8°C. Partially used vials should be discarded.

Do not freeze. Before use, visually inspect the medicine. The solution must be clear or slightly opalescent and colorless or pale yellow. Do not use if the solution is cloudy or has deposits.

Keep out of reach and sight of children. Store in the original container to protect from light.

DETAILS OF PERMISSION: Manufacturing License number: G28E/1

Report suspected adverse reaction at: Hemofluidsafety@intaspharma.com

DATE OF PREPARATION: 27.01.2022

Manufactured and Marketed by: (NTAS)

INTAS PHARMACEUTICALS LTD.

Plot No. 4961/A&B, Sarkhej-Bavla Highway,

Village: Matoda, Taluka: Sanand,

Ahmedabad-382213, Gujarat, INDIA.

AW-137-00

Front Side

Back Side

Reason for revision : As per change control No. CBL/CCF/2021/0237

Artwork No.: AW-137-00

Supersedes No.: NA

Approval table with columns: Prepared by, Reviewed by, Approved by, Department, Sign, Date, Name, Designation. Rows include QA, Production, Regulatory, Marketing, Medical, and QA.

Artwork Name : GlobuceL VF-10 (IP) - leaflet
Size : 130 x 290 mm
GSM of Paper : 50 ± 10% GSM

Colour Shade No. : Black
No. of fold : 4
Fold size : ~65 mm X ~36.25 mm

Type of Paper : Maplitho
27.01.2022